

FeynRules

Beyond the Standard Model phenomenology made easy.

Benjamin Fuks (IPHC Strasbourg)

In collaboration with N. Christensen (MSU), C. Duhr (UCL), FeynRules people & MadGraph people.

GDR Terascale
LPSC Grenoble
March 31, 2009

Outline

- 1 Motivation: a roadmap to BSM at the LHC
- 2 FeynRules
- 3 Model database and validation status
- 4 Summary - outlook

Outline

- 1 Motivation: a roadmap to BSM at the LHC
- 2 FeynRules
- 3 Model database and validation status
- 4 Summary - outlook

A roadmap to BSM at the LHC (1)

Models

Theoretical works

- * **Pen&pencil stage.**
- * Leading order, loop calculations, ...
- * Electroweak, low energy constraints,...

Phenomenological works

- * **Monte Carlo event generation.**
⇒ **Feynman rules tables!**
- * Generic detector simulation, ...
- * Signal/background studies.

Experimental works


- * **Validated experimental framework.**
⇒ **Contains Monte Carlo generators!**
- * Realistic detector simulation, ...
- * Comparison with data.




Data

How to go from models to data?

A roadmap to BSM at the LHC (2)



A roadmap to BSM at the LHC (3)



A roadmap to BSM at the LHC (4)

Models

F
E
Y
N
R
U
L
E
S



Theoretical works

- * **Pen&pencil stage.**
- * Leading order, loop calculations, ...
- * Electroweak, low energy constraints, ...



Phenomenological works

- * **Monte Carlo event generation.**
⇒ **Feynman rules tables!**
- * Generic detector simulation, ...
- * Signal/background studies.



Experimental works

- * **Validated experimental framework.**
⇒ **Contains Monte Carlo generators!**
- * Realistic detector simulation, ...
- * Comparison with data.



Data


Solution 3: FeynRules.

- Communicates with MadGraph, Sherpa, (Pythia, Herwig), ...
- ✓ No new MC validation.
- ✓ MC validated for exp. software.
- ✓ Mathematica based.
- ✓ Portable.

Outline

- 1 Motivation: a roadmap to BSM at the LHC
- 2 FeynRules**
- 3 Model database and validation status
- 4 Summary - outlook

FeynRules



Example: QCD - Parameters

Parameters of the model

```
aS == {
  Description      -> "Strong coupling constant at MZ"
  Tex              -> Subscript[\[Alpha], s],
  ParameterType    -> External,
  BlockName        -> SMINPUTS,
  OrderBlock       -> 3,
  InteractionOrder -> {QCD, 2}},
gs == {
  Description      -> "Strong coupling constant",
  Tex              -> Subscript[g, s],
  ComplexParameter -> False,
  ParameterType    -> Internal,
  Value            -> Sqrt[4 Pi aS],
  InteractionOrder -> {QCD, 1},
  ParameterName    -> "G"}
```

- * **All the information** needed by the MC codes.
- * **TeX-form** (for the TeX-file).
- * **Complex/real** parameters.
- * **External/internal** parameters.

Example: QCD - Gauge group and gauge boson

The $SU(3)_C$ gauge group

```
SU3C == {  
  Abelian           -> False,  
  GaugeBoson       -> G,  
  StructureConstant -> f,  
  DTerm            -> dSUN,  
  Representations   -> {T, Colour},  
  CouplingConstant -> gs}
```

Gluon field definition

```
V[1] == {  
  ClassName       -> G,  
  SelfConjugate   -> True,  
  Indices         -> Index[Gluon],  
  Mass            -> 0,  
  Width           -> 0,  
  ParticleName    -> "g",  
  PDG             -> 21,  
  PropagatorLabel -> "G",  
  PropagatorType  -> C,  
  PropagatorArrow -> None}
```

- * **Gauge boson** definition.
- * **Gauge group** definition.
- * Association of a **coupling constant**.
- * Definition of the **structure functions**.
- * Definition of the **representations**.

Example: QCD - Quark fields

The quark fields

```
F[1] == {  
  ClassName      -> q,  
  ClassMembers   -> {d, u, s, c, b, t},  
  FlavorIndex    -> Flavour,  
  SelfConjugate  -> False,  
  Indices        -> {Index[Flavour], Index[Colour]},  
  Mass           -> {MQ, MD, MU, MS, MC, MB, MT},  
  Width         -> {WQ, 0, 0, 0, 0, 0, WT},  
  ParticleName   -> {"d", "u", "s", "c", "b", "t"},  
  AntiParticleName -> {"d~", "u~", "s~", "c~", "b~", "t~"},  
  PDG           -> {1, 2, 3, 4, 5, 6},  
  PropagatorLabel -> {"q", "d", "u", "s", "c", "b", "t"},  
  PropagatorType -> Straight,  
  PropagatorArrow -> Forward}
```

- * **Classes:** implicit sums in the Lagrangian.
- * **All the information** needed by the MC codes.

Example: QCD - Lagrangian

The QCD Lagrangian

```
LQCD = -1/4 * FS[G, mu, nu, a] * FS[G, mu, nu, a] +
I*qbar.Ga[mu].del[q, mu] - MQ[f] * qbar[s,f,c].q[s,f,c] +
gs * G[mu,a] * qbar.Ga[mu].T[a].q
```

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu} + \sum_f \left[\bar{q}_f (i\not{\partial} - m_f + g_s \mathbb{G}^a T^a) q_f \right],$$

where we are summing over the quark flavours.

- * **Gluon strength tensor**: automatically defined with the gauge group.
- * **Implicit summations** \Rightarrow easy debugging.

Example: QCD - Results

Results - let us do (some) phenomenology!

```
FeynmanRules[LQCD, FlavorExpand->False]
```

```
Vertex 1
```

```
Particle 1 : Vector , G
```

```
Particle 2 : Dirac , q†
```

```
Particle 3 : Dirac , q
```

```
Vertex:
```

$$i g_s \gamma_{s_2, s_3}^{\mu_1} \delta_{f_2, f_3} T_{m_2, m_3}^a$$

```
WriteFeynArtsOutput[LQCD]
```

```
WriteCHOutput[LQCD]
```

```
WriteMGOutput[LQCD]
```

```
WriteSHOutput[LQCD]
```

Explicit **flavour expansion**: six vertices instead of one.

Outline

- 1 Motivation: a roadmap to BSM at the LHC
- 2 FeynRules
- 3 Model database and validation status**
- 4 Summary - outlook

Model database

- **Publicly available (FeynRules v1.2.5):**
 - * **The Standard Model (SM)** [N. Christensen, C. Duhr].
 - * **Higgs effective theory (large m_{top} approximation)** [C. Duhr].
 - * **The Three-Site Model** [N. Christensen].
 - 5D $SU(2) \times SU(2) \times U(1)$ theory in a slice of Anti-deSitter space.
 - Gauge invariant higgsless model.
 - Heavy (≈ 400 GeV) and nearly degenerate extra gauge bosons.
 - SM-like plus new fermionic states.
 - * **The Hill Model** [P. Aquino, C. Duhr].
 - SM plus an additional scalar sector coupling only to the Higgs.
 - Two Higgs fields after mass matrix diagonalization.
- **Soon publicly available:**
 - * **The general MSSM** (with 105 free parameters) [BenjF].
 - * **Minimal universal extra dimensions** [P. Aquino].
 - * **Effective QCD for the pseudoscalar nonet** [C. Degrande].
 - * **Four generations models** with right-handed neutrinos [BenjF, M. Spannowsky].
 - * **Left-right symmetric models** [L. Basso].
 - * **Little Higgs model** [T. Figy].
 - * **Effective quantum gravity** [BenjF, C. Reuschle].
 - * **Type-III See-Saw model** [R. Franceschini].

Validation sheet

- **FeynArts/FormCalc:**
 - * Use of the **FeynRules** version of the FeynArts model files.
 - * **Check of the FormCalc-produced formulas with literature.**
 - * Used versions: **FormCalc-5.4** and **FormCalc-6.0**.
- **MadGraph/MadEvent:**
 - * Comparison between (existing) **stock** and **FeynRules** model files.
 - * Test of various $2 \rightarrow 2$ and $2 \rightarrow 3$ processes.
 - * Used version: **MadGraph-4.4.17**.
- **CalcHEP/CompHEP:**
 - * Comparison between (existing) **stock** and **FeynRules** model files.
 - * **Test of both Feynman and unitary gauges.**
 - * Test of various $2 \rightarrow 2$ and $2 \rightarrow 3$ processes.
 - * Used version: **CalcHEP-2.5**.
- **Sherpa:**
 - * To be done...

Example: validation status of the MSSM (1)

- **Handmade vs. automated implementation.**
 - * 2522 vertices, without the four-scalar interactions.
 - * **More than 10000 vertices, with the four-scalar interactions !!!**
- **FeynArts/FormCalc:** ongoing...
 - ✓ **FormCalc-5.4:** all $2 \rightarrow 2$ **SUSY** particle pair hadroproduction processes.
 - ✓ **FormCalc-6.0:** in the flavour conserving MSSM (cMSSM) limit.
 - ➔ **FormCalc-6.0:** almost there in the general MSSM.
- ✓ **MadGraph/MadEvent** (in the cMSSM limit):
 - * MG-Stock was validated by the CATPISS collaboration [Hagiwara *et al.* (2006)].
 - ✓ **320 decay widths.**
 - ✓ **456 $2 \rightarrow 2$ SUSY processes.**
 - ✓ **2708 $2 \rightarrow 3$ SUSY processes.**

The signs and absolute values of all the vertices have been checked.

- **CalcHEP/CompHEP (in the cMSSM):** ongoing check for $2 \rightarrow 2$ processes.
 - * **Some bugs found in the stock version!**

Example: validation status of the MSSM (2)

Some MadGraph and CalcHEP results

Process	MG-FR	MG-stock	CH-FR	CH-Stock	Result
e+,e->e+,e-	7.5203×10^{-2}	7.5216×10^{-2}	7.5137×10^{-2}	7.5137×10^{-2}	OK: 0.105086%
e+,e->vm,vm-	1.5268×10^{-3}	1.5285×10^{-3}	1.5261×10^{-3}	1.5262×10^{-3}	OK: 0.15714%
e+,e->t,t-	1.1098×10^{-2}	1.1101×10^{-2}	1.1108×10^{-2}	1.1114×10^{-2}	OK: 0.144066%
e+,e->d,d-	5.6391×10^{-3}	5.6597×10^{-3}	5.6465×10^{-3}	5.6465×10^{-3}	OK: 0.36464%
e+,e->W+,W-	2.8014×10^{-1}	2.801×10^{-1}	2.8008×10^{-1}	2.8009×10^{-1}	OK: 0.0214202%
e+,e->Z,Z	1.535×10^{-2}	1.5347×10^{-2}	1.5347×10^{-2}	1.5347×10^{-2}	OK: 0.0195459%
e+,e->Z,a	6.2902×10^{-2}	6.2901×10^{-2}	6.292×10^{-2}	6.292×10^{-2}	OK: 0.0302016%
e+,e->s15-,s15+	3.2044×10^{-2}	3.2002×10^{-2}	3.2039×10^{-2}	3.2039×10^{-2}	OK: 0.131156%
e+,e->s12-,s12+	3.6401×10^{-2}	3.641×10^{-2}	3.64×10^{-2}	3.64×10^{-2}	OK: 0.0274688%
e+,e->s15-,s12+	2.0292×10^{-3}	2.0269×10^{-3}	2.0291×10^{-3}	2.0291×10^{-3}	OK: 0.113409%
e+,e->s11-,s11+	1.6061×10^{-3}	1.6061×10^{-3}	1.6054×10^{-3}	1.6054×10^{-3}	OK: 0.0435933%
e+,e->sv3,sv3-	9.5578×10^{-2}	9.5567×10^{-2}	9.554×10^{-2}	9.5542×10^{-2}	OK: 0.039766%
e+,e->su4,su4-	2.9679×10^{-3}	2.9676×10^{-3}	2.9692×10^{-3}	2.9692×10^{-3}	OK: 0.0539011%
e+,e->su1,su1-	1.9518×10^{-3}	1.9486×10^{-3}	1.9517×10^{-3}	1.9517×10^{-3}	OK: 0.164086%
e+,e->su6,su6-	2.2021×10^{-3}	2.2041×10^{-3}	2.202×10^{-3}	2.202×10^{-3}	OK: 0.0953224%
e+,e->su1,su6-	4.4196×10^{-4}	4.4134×10^{-4}	4.4155×10^{-4}	4.4155×10^{-4}	OK: 0.140383%
e+,e->sd4,sd4-	4.9197×10^{-4}	4.926×10^{-4}	4.9192×10^{-4}	4.9192×10^{-4}	OK: 0.138138%
e+,e->sd6,sd6-	2.0014×10^{-3}	2.0012×10^{-3}	2.0016×10^{-3}	2.0016×10^{-3}	OK: 0.019986%
e+,e->sd1,sd2-	2.1502×10^{-4}	2.149×10^{-4}	2.1494×10^{-4}	2.1494×10^{-4}	OK: 0.0558243%
e+,e->n1,n1	7.6112×10^{-3}	7.6075×10^{-3}	7.6077×10^{-3}	7.6076×10^{-3}	OK: 0.0486244%
e+,e->n1,n3	2.7949×10^{-3}	2.792×10^{-3}	2.7942×10^{-3}	2.7943×10^{-3}	OK: 0.103814%
e+,e->n2,n2	4.1779×10^{-4}	4.1709×10^{-4}	4.17×10^{-4}	4.1701×10^{-4}	OK: 0.189269%
e+,e->n2,n4	7.5931×10^{-4}	7.5959×10^{-4}	7.5912×10^{-4}	7.5914×10^{-4}	OK: 0.0618946%
e+,e->n4,n4	3.5319×10^{-5}	3.531×10^{-5}	3.5317×10^{-5}	3.5317×10^{-5}	OK: 0.0254853%
e+,e->x1+,x1-	1.204×10^{-2}	1.2038×10^{-2}	1.2039×10^{-2}	1.2039×10^{-2}	OK: 0.0166127%
e+,e->x2+,x2-	7.0411×10^{-3}	7.0479×10^{-3}	7.0494×10^{-3}	7.0494×10^{-3}	OK: 0.11781%
e+,e->Z,h1	7.6379×10^{-4}	7.6496×10^{-4}	7.6477×10^{-4}	7.6478×10^{-4}	OK: 0.153066%
e+,e->Z,h2	1.0024×10^{-7}	1.0007×10^{-7}	1.0017×10^{-7}	1.0017×10^{-7}	OK: 0.169737%
e+,e->h3,h1	9.9472×10^{-8}	9.9485×10^{-8}	9.9461×10^{-8}	9.9466×10^{-8}	OK: 0.0241272%
e+,e->h3,h2	7.172×10^{-4}	7.1771×10^{-4}	7.177×10^{-4}	7.1771×10^{-4}	OK: 0.0710846%
e+,e->H+,H-	1.7338×10^{-3}	1.7338×10^{-3}	1.7355×10^{-3}	1.7355×10^{-3}	OK: 0.0980025%

Example: validation status of minimal UED model

- ✓ **MadGraph/MadEvent:**
 - * No MG-Stock.
 - ✓ **118 2 → 2 processes.**
- ✓ **CalcHEP/CompHEP: 118 2 → 2 processes.**
 - * Stock version: Datta, Kong and Matchev implementation.
 - * **One bug found in the stock version!**

Some results

Process	MG-FR	CH-FR	CH-Stock	Result
e1R-,e1R->u,u-	1.107×10^{-1}	1.1094×10^{-1}	1.1094×10^{-1}	OK: 0.216567%
e1R-,e1R->d,d-	3.277×10^{-2}	3.2795×10^{-2}	3.2795×10^{-2}	OK: 0.0762602%
e1R-,e1R->e-,e+	2.5553×10^{-1}	2.5537×10^{-1}	2.5537×10^{-1}	OK: 0.0626346%
e1R-,e1R->e-,e-	1.0714	1.0714	1.0714	OK: 0.0%
e1R-,n1R->e-,n-	6.5807×10^{-1}	6.5818×10^{-1}	6.5818×10^{-1}	OK: 0.0167142%
e1R-,n1R->e-,n+	4.7857×10^{-1}	4.7682×10^{-1}	4.7682×10^{-1}	OK: 0.366343%
e1R-,e1R->A,A	2.0803×10^{-1}	2.0788×10^{-1}	2.0788×10^{-1}	OK: 0.072131%
n1l,n1l->u,u-	1.6364×10^{-1}	1.6354×10^{-1}	1.6354×10^{-1}	OK: 0.0611284%
n1l,n1l->Z,Z	4.1402×10^{-1}	4.1349×10^{-1}	4.1349×10^{-1}	OK: 0.128095%
n1l,n1l->W+,W-	5.9018×10^{-1}	5.9009×10^{-1}	5.901×10^{-1}	OK: 0.0152507%
e1L-,e1L->u,u-	2.3023×10^{-1}	2.2977×10^{-1}	2.2977×10^{-1}	OK: 0.2%
e1L-,e1L->d,d-	1.4289×10^{-1}	1.4274×10^{-1}	1.4275×10^{-1}	OK: 0.105031%
e1L-,e1L->e-,e+	2.5×10^{-1}	2.4978×10^{-1}	2.4978×10^{-1}	OK: 0.0880387%
e1L-,n1l->d,u-	6.3986×10^{-1}	6.3998×10^{-1}	6.3999×10^{-1}	OK: 0.0203149%
e1L-,n1l->e-,n1-	6.3118×10^{-1}	6.3132×10^{-1}	6.3133×10^{-1}	OK: 0.0237622%
e1L-,n1l>e-,n1	1.0519	1.0519	1.0519	OK: 0.0%
B1,B1>u,u-	9.2638×10^{-2}	9.2548×10^{-2}	9.2548×10^{-2}	OK: 0.0971996%
B1,B1>d,d-	6.1392×10^{-3}	6.1347×10^{-3}	6.1347×10^{-3}	OK: 0.0733263%
B1,B1>e+,e-	1.8444×10^{-1}	1.8411×10^{-1}	1.8411×10^{-1}	OK: 0.17908%
Z1,Z1>u,u-	3.5574×10^{-1}	3.5556×10^{-1}	3.5556×10^{-1}	OK: 0.0506116%
Z1,Z1>d,d-	3.566×10^{-1}	3.5556×10^{-1}	3.5556×10^{-1}	OK: 0.292069%
Z1,Z1>e+,e-	1.3429×10^{-1}	1.3409×10^{-1}	1.3409×10^{-1}	OK: 0.149042%
Z1,Z1>W-,W+	2.8571×10^1	2.8573×10^1	2.8573×10^1	OK: 0.00699986%

Example: validation status of the Three-Site model

✓ MadGraph/MadEvent:

* No MG-Stock.

✓ 224 2 → 2 processes.

✓ CalcHEP/CompHEP: 224 2 → 2 processes.

* Stock version: N. Christensen's implementation.

Some results

Process	MG-FR	CH-FR	CH-Stock	Result
-W _s ,W->W _s ,W-	3.8293×10^1	3.8243×10^1	3.8218×10^1	OK: 0.19605%
-W _s ,W->W _s ,W-	3.8365×10^1	3.8244×10^1	3.8219×10^1	OK: 0.381281%
-W _s ,W->Z,Z	4.6001×10^1	4.5986×10^1	4.5956×10^1	OK: 0.0978718%
-W _s ,W->Z,Z	4.5986×10^1	4.5986×10^1	4.5956×10^1	OK: 0.0652585%
-W _s ,W->A,Z	1.6925×10^1	1.6891×10^1	1.688×10^1	OK: 0.266233%
-W _s ,W->A,Z	1.684×10^1	1.6891×10^1	1.688×10^1	OK: 0.302392%
-Z,Z->W _s ,W-	8.2375×10^1	8.2402×10^1	8.2349×10^1	OK: 0.0643395%
-Z,Z->W _s ,W-	6.8543×10^1	6.871×10^1	6.8666×10^1	OK: 0.243346%
-W _s ,W->A,A	6.0926×10^{-18}	-1.2302×10^{-31}	3.1165×10^{-32}	Discrepancy: 200.%
-W _s ,W->A,A	6.098×10^{-18}	-1.2302×10^{-31}	3.1165×10^{-32}	Discrepancy: 200.%
Process	MG-FR	CH-FR	CH-Stock	Result
u1,-D1->W _s ,G	6.4153×10^{-2}	6.4112×10^{-2}	6.4091×10^{-2}	OK: 0.0966907%
u1,-D1->W _s ,-Z	3.8531×10^{-1}	3.8539×10^{-1}	3.8514×10^{-1}	OK: 0.0648904%
u1,-D1->W _s ,-Z	4.3576×10^{-1}	4.3594×10^{-1}	4.3566×10^{-1}	OK: 0.0642497%
-u1,-D1->W _s ,G	1.0137	1.0139	1.0136	OK: 0.0295931%
-u1,-D1->W _s ,G	1.1532	1.1523	1.152	OK: 0.104112%
-u1,-D1->W _s ,-Z	2.8898×10^{-1}	2.8845×10^{-1}	2.8826×10^{-1}	OK: 0.249463%
u1,-D1->W _s ,G	1.0173	1.0139	1.0136	OK: 0.36437%
u1,-D1->W _s ,G	1.1543	1.1523	1.1519	OK: 0.208135%
u1,-D1->W _s ,-Z	2.9225×10^{-1}	2.9273×10^{-1}	2.9254×10^{-1}	OK: 0.164108%
-u1,-D1->W _s ,G	1.5625×10^{-1}	1.5605×10^{-1}	1.56×10^{-1}	OK: 0.160128%
-u1,-D1->W _s ,G	1.0343	1.0358	1.0355	OK: 0.144921%
-u1,-D1->W _s ,-Z	2.7826	2.7837	2.7819	OK: 0.0646831%
-u1,-D1->W _s ,-Z	2.5678	2.5706	2.5689	OK: 0.108983%

Outline

- 1 Motivation: a roadmap to BSM at the LHC
- 2 FeynRules
- 3 Model database and validation status
- 4 Summary - outlook**

Summary: the philosophy of FeynRules

- * **Theorist-friendly environment** to develop new models.
Mathematica-based.
- * **Filling the gap** between model building and collider phenomenology.
 - 1) Lagrangian → FeynRules → model files for your favourite Monte Carlo codes.
 - 2) Monte Carlo code → phenomenology.
- * **Avoid separate implementations** of a model on different programs.
FeynRules does it for you!
- * **Exploit the strengths of the different programs!**
- * **The validation of the existing models is ongoing.**

- * Contact us to add your favourite **model**.
- * Contact us to add your favourite **Monte Carlo tool**.
- * Website: <http://feynrules.phys.ucl.ac.be> .